


INTRODUCTION

Classical muscle biopsy studies¹ demonstrated that elite **endurance** athletes exhibit a more pronounced **slow** muscle typology, while elite athletes who excel in **explosive** disciplines, have a predominant **fast** muscle typology. **Muscle carnosine quantification** by proton magnetic resonance spectroscopy (¹H-MRS) was developed as a **non-invasive alternative to estimate muscle typology**². The validity of this technique was initially demonstrated in track-and-field, where muscle carnosine levels in elite athletes gradually decrease with increasing running distance. Also in **cycling**, it is anticipated that the various disciplines span a range of muscle typologies, but **solid data are scarce**.



Proton Magnetic Resonance Spectroscopy (¹H-MRS)
'Muscle Talent Scan'

AIMS

- 1 To compare the **muscle carnosine levels** of elite cyclists, excelling in **different cycling disciplines**.
- 2 To investigate **transfer possibilities** between cycling disciplines, **based on muscle typology**.

METHODS

Muscle carnosine levels of **85 elite cyclists** excelling in 8 different cycling disciplines were measured with the Muscle Talent Scan (¹H-MRS) in the 2 calf muscles: m. soleus and m. gastrocnemius.

Elite cyclist
European/World Championship participant
UCI WorldTour Team
Pro Continental Cycling Team



5 multi stage riders



14 single stage riders



5 road sprinters



4 BMX riders



29 track endurance cyclists



9 track-sprinters




11 mountain bikers



9 cyclo-crossers

To investigate the muscle typology of the cyclists, their muscle carnosine levels were converted to z-scores, based on the normal carnosine distribution of a control population (112 women, 163 men).




RESULTS

1 Carnosine levels


Muscle typology distribution for all cycling disciplines.

Slow Intermediate Fast



2 Talent transfer

Differences (p-values) in mean carnosine levels between the 8 cycling disciplines. The more the disciplines differ from each other, the less transfer seems possible, based on muscle typology.



CONCLUSIONS

- 1 Prominent differences in muscle carnosine levels exist between elite cyclists of various disciplines. Road cyclists, cyclo-crossers and mountain bikers display low carnosine levels, indicative for a slow muscle typology, whereas the carnosine levels of BMX riders and track-sprinters are indicative for a dominant fast muscle typology.
- 2 The non-invasive character of the Muscle Talent Scan opens opportunities for application in transfer and talent orientation in cycling.

REFERENCES

Costill, D.L., Daniels, J., Evans, W., Fink, W., Krahenbuhl, G., and Saltin, B. (1976). Skeletal muscle enzymes and fiber composition in male and female track athletes. J. Appl. Physiol. 40, 149–154.
Gollnick, P.D., Armstrong, R.B., Saubert IV, C.W., Piehl, K., and Saltin, B. (1972). Enzyme activity and fiber composition in skeletal muscle of untrained and trained men. J. Appl. Physiol. 33, 312–319.
Baguet, A., Everaert, I., Hespel, P., Petrovic, M., Achten, E., and Derave, W. (2011a). A new method for non-invasive estimation of human muscle fiber type composition. PLoS One 6, 1–6.



@TeamDerave



www.musclelalentalscan.com



Kim.VanVossel@ugent.be



info@musclelalentalscan.com